

**OXFORD CAMBRIDGE AND RSA EXAMINATIONS**

**Advanced GCE**

**CHEMISTRY**

**2815/06**

**Transition Elements**

**Tuesday**

**25 JANUARY 2005**

**Afternoon**

**50 minutes**

Candidates answer on the question paper.

Additional materials:

*Data Sheet for Chemistry*

Scientific calculator

Candidate Name	Centre Number	Candidate Number												
	<table border="1" style="display: inline-table; border-collapse: collapse;"> <tr> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> </tr> </table>							<table border="1" style="display: inline-table; border-collapse: collapse;"> <tr> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> </tr> </table>						

**TIME** 50 minutes

**INSTRUCTIONS TO CANDIDATES**

- Write your name in the space above.
- Write your Centre number and Candidate number in the boxes above.
- Answer **all** the questions.
- Write your answers in the spaces provided on the question paper.
- Read each question carefully and make sure you know what you have to do before starting your answer.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- You will be awarded marks for the quality of written communication where this is indicated in the question.
- You may use a scientific calculator.
- You may use the *Data Sheet for Chemistry*.
- You are advised to show all the steps in any calculations.

FOR EXAMINER'S USE		
Qu.	Max.	Mark
1	12	
2	11	
3	13	
4	9	
<b>TOTAL</b>	<b>45</b>	

**This question paper consists of 10 printed pages and 2 blank pages.**

Answer all the questions.

- 1 Vanadium is a hard corrosion-resistant metal which forms compounds with a number of different oxidation states.

(a) State a common use for vanadium or one of its compounds.

..... [1]

(b) The standard electrode potential of the  $V^{2+}/V$  redox system is  $-1.20\text{ V}$ .

Draw a labelled diagram to show how you would measure the standard electrode potential of the  $V^{2+}/V$  system.

[5]

(c) The most common oxidation states of vanadium are shown in the table below.

	$V^{2+}$	$VO_2^+$	$VO^{2+}$	$V^{3+}$
oxidation number of vanadium	+2			+3
colour	lilac	yellow		




(i) Complete the table by filling in the empty spaces.

[4]

- (ii) Each oxidation state may be observed by carrying out the successive reduction of ammonium vanadate(V) using zinc in an acidic solution.

The final step converts  $V^{3+}(aq)$  into  $V^{2+}(aq)$ .

Use the following standard electrode potentials to explain why the reduction process stops at the ion  $V^{2+}$ .



.....

.....

.....

..... [2]

[Total: 12]

2 Transition metals readily form complex ions when they are combined with a suitable ligand.

(a) What is meant by the following terms?

(i) *complex ion*

.....  
..... [1]

(ii) *ligand*

.....  
..... [2]

(b) A common ligand which combines with a number of transition metal ions is ethane-1,2-diamine,  $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$ . This is a bidentate ligand.

Explain the meaning of the term *bidentate*.


.....  
..... [1]



3 One common property of transition metal compounds is that they are coloured. When a transition metal ion forms a complex, splitting of the d-orbitals takes place.

(a) In a  $\text{Cu}^{2+}$  ion, all five d-orbitals have the same energy. However, when the octahedral complex ion  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$  is formed, the d-orbitals split into different energy levels.

(i) Complete the following diagram to show the splitting of d-orbitals in the complex ion.




$\text{Cu}^{2+}$


$[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$

[2]

(ii) Draw diagrams to show the shape of one lower energy d-orbital and one higher energy d-orbital in the boxes below.



lower energy  
d-orbital




higher energy  
d-orbital

[2]



- (c) If visible light is passed through a coloured solution, the light that is transmitted can be analysed by a visible spectrometer.

The visible spectrum for aqueous  $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$  is shown below.



The solution is purple. Explain how you can tell that it is purple by looking at the spectrum.

.....


.....

..... [2]

[Total: 13]



- 4 Potassium dichromate(VI) can be used in a number of redox reactions. The standard electrode potentials for two half reactions are given below.



- (a) Acidified potassium dichromate(VI) is added to aqueous potassium iodide to give aqueous iodine.


- (i) Construct an ionic equation to show the reaction taking place when acidified potassium dichromate(VI) is added to aqueous potassium iodide.

.....  
 .....  
 ..... [2]

- (ii) An excess of aqueous sodium thiosulphate was then added. Describe and explain what you would see.

.....  
 .....  
 .....  
 ..... [3]

- (b) Potassium dichromate(VI) also takes part in the following reaction.



- (i) Show that chromium is **not** taking part in a redox reaction.

.....  
 .....  
 ..... [2]

- (ii) Describe the colour change for the forward reaction.

from ..... to ..... [1]

- (iii) Suggest a reagent that would convert  $\text{CrO}_4^{2-}$  back to  $\text{Cr}_2\text{O}_7^{2-}$ .

..... [1]

[Total: 9]

**END OF QUESTION PAPER**