

OXFORD CAMBRIDGE AND RSA EXAMINATIONS

Advanced GCE

CHEMISTRY

2816/01

Unifying Concepts in Chemistry

Friday

24 JANUARY 2003

Afternoon

1 hour 15 minutes

Candidates answer on the question paper.

Additional materials:

Data Sheet for Chemistry

Scientific calculator

Candidate Name	Centre Number	Candidate Number												
	<table border="1" style="display: inline-table; border-collapse: collapse;"> <tr> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> </tr> </table>							<table border="1" style="display: inline-table; border-collapse: collapse;"> <tr> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> <td style="width: 15px; height: 15px;"></td> </tr> </table>						

TIME 1 hour 15 minutes

INSTRUCTIONS TO CANDIDATES

- Write your name in the space above.
- Write your Centre number and Candidate number in the boxes above.
- Answer **all** the questions.
- Write your answers in the spaces provided on the question paper.
- Read each question carefully and make sure you know what you have to do before starting your answer.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- You will be awarded marks for the quality of written communication where this is indicated in the question.
- You may use a scientific calculator.
- You may use the *Data Sheet for Chemistry*.
- You are advised to show all the steps in any calculations.


FOR EXAMINER'S USE		
Qu.	Max.	Mark
1	17	
2	17	
3	12	
4	14	
TOTAL	60	

This question paper consists of 10 printed pages and 2 blank pages.

Answer all the questions.

- 1 *Syngas* is a mixture of carbon monoxide and hydrogen gases, used as a feedstock for the manufacture of methanol.

A dynamic equilibrium was set up between carbon monoxide, CO, hydrogen, H₂, and methanol, CH₃OH. The equilibrium system is shown by Equilibrium 1.1 below.



The equilibrium concentrations of the three components of this equilibrium are shown below.

component	CO(g)	H ₂ (g)	CH ₃ OH(g)
equilibrium concentration /mol dm ⁻³	3.1 × 10 ⁻³	2.4 × 10 ⁻²	2.6 × 10 ⁻⁵

- (a) State two features of a system that is in *dynamic equilibrium*.

.....

 [2]

- (b) (i) Write the expression for K_c for this equilibrium system.

[2]

- (ii) Calculate the numerical value of K_c for this equilibrium.

[2]

- (c) The pressure was increased whilst keeping the temperature constant. The system was left to reach equilibrium. The equilibrium position of Equilibrium 1.1 shifted to the right.

- (i) Explain why the equilibrium moved to the right.

.....
 [2]

- (ii) What is the effect, if any, on K_c ?

..... [1]

(iii) State and explain the effect on the rates of the forward and reverse reactions

- when the pressure was first changed
- when the system reached equilibrium.

.....
.....
.....
.....
.....[4]

(d) The temperature was increased whilst keeping the pressure constant. The system was left to reach equilibrium. The value of K_c for Equilibrium 1.1 decreased.

(i) Explain what happens to the equilibrium position of Equilibrium 1.1.

.....
.....
.....
.....
.....[2]

(ii) Deduce the sign of the enthalpy change for the forward reaction shown in Equilibrium 1.1. Explain your reasoning.

.....
.....[1]

(iii) Explain how the partial pressure of $\text{CH}_3\text{OH}(\text{g})$ would change as the system moves towards equilibrium.

.....
.....
.....[1]

[Total: 17]

- 2 Nitrous oxide, N_2O , is a colourless gas with a mild, pleasing odour and sweet taste. It is widely used as a propellant in aerosol cans of whipped cream.

(a) Nitrous oxide is formed when ammonium nitrate, NH_4NO_3 , is gently heated.



(i) What mass of N_2O is formed by heating 100 g of NH_4NO_3 ?


[3]

(ii) What happens to the oxidation number of each nitrogen from NH_4NO_3 in this reaction?

.....


[2]

(b) When heated strongly, nitrous oxide decomposes into its elements.



This reaction is first order with respect to N_2O .

The graph below shows how nitrous oxide decomposes with time at constant temperature.



(i) Explain how the graph confirms that this reaction is first order with respect to N₂O.

.....
.....
.....
.....[3]

(ii) Write the expression for the rate equation of this reaction.

[1]

(iii) Use the graph to work out the rate of reaction, in mol dm⁻³ s⁻¹, at 70 seconds. Show clearly your working on the graph.

rate =mol dm⁻³ s⁻¹ [2]

(iv) Calculate the rate constant for this reaction. State the units.

k = units [2]

(v) What evidence is there that the mechanism of this reaction takes place in more than a single step?

.....
.....
.....[2]

(c) N₂O is occasionally injected into the engines of racing cars to give more power and exceptional acceleration. The N₂O decomposes exothermically to N₂ and O₂.

Suggest **two** reasons why this reaction provides an extra boost to the engine.

.....
.....
.....[2]

[Total: 17]

(b) The acid dissociation constant K_a of hydrocyanic acid, HCN, is $4.9 \times 10^{-10} \text{ mol dm}^{-3}$.

(i) Write an expression for the acid dissociation constant of HCN.

[1]

(ii) Calculate the pH of a $0.010 \text{ mol dm}^{-3}$ solution of hydrocyanic acid.

[3]

[Total: 12]

4 Organic acids occur widely in nature.

- (a) Butanoic acid, $\text{CH}_3(\text{CH}_2)_2\text{COOH}$, is a straight-chain organic acid, largely responsible for the odour of rancid butter.

Caprylic acid is another straight-chain organic acid. It is produced in the body in small amounts as an antifungal agent in human sweat.

- (i) Some caprylic acid was isolated from human sweat and analysed. The sample of caprylic acid had the percentage composition by mass:

C, 66.7%; H, 11.1%; O, 22.2%. $M_r = 144$.

Calculate the molecular formula of caprylic acid and suggest its structural formula.

[4]

- (ii) Tracker dogs are trained to follow odours such as the characteristic blend of organic acids in the sweat from a person's feet. A dog is able to detect extremely small quantities of these acids.

Sweat containing equal amounts of butanoic and caprylic acids produces more butanoic acid vapour than caprylic acid vapour.

Suggest a reason for this. Explain your answer.

.....
.....
.....
.....[2]

