

## 2.2 Group 2 – The alkaline earth metals

**Atomic radius – Increases down the Group:**

- **Shells:** **More** electron shells
- **Shielding:** **More** shielding (more inner shells)

**Ionic radius – Smaller** than atomic radius:

- **1 less shell:** **As 2e lost removing outer shell**

**First ionisation - Decreases down the Group:**

Number of shells increases

Shielding increases

Atomic radius increases



First ionisation energy decreases

- **Shells:** **More** electron shells
- **Shielding:** **More** shielding (more inner shells)
- **No. Protons:** Number of protons **increases** but is **outweighed** by shells and shielding
- **Attraction:** Therefore attraction is **less**
- **Energy:** **Energy** required to remove an electron **decreases**

**Reactivity – Increases down the Group:**

- All lose 2 electrons forming a 2+ ion when they react
- **Ionisation energies decrease** as you go down the group
- **Electrons are lost more easily**
- **Reactivity increases** as you go down the group

**Melting point – decreases down the group:**




- As you go down Group 2 the **ionic radius increases**
- The 2+ charge from the nucleus is **further away** from the delocalised electrons
- **Attraction** is therefore **weaker**
- **Energy** required is **less**

- Mg's unusually low melting point comes from the different arrangement of the ions in the crystal structure.

### Reactions of the Group 2 elements:



#### How the Group 2 elements react:

- Group 2 metals are reactive and all **lose 2e** when they react.
- As you go down Group 2 they become **more reactive**.
- This is due to the **decrease in ionisation energies** as you go down the group.



#### Reaction with water

- Group 2 metals react with water to give the hydroxide and hydrogen gas:



### Solubility of the hydroxides and sulphates:

Mg(OH) <sub>2</sub>		Least soluble		MgSO <sub>4</sub>		Most soluble
Ca(OH) <sub>2</sub>				CaSO <sub>4</sub>		
Sr(OH) <sub>2</sub>				SrSO <sub>4</sub>		
Ba(OH) <sub>2</sub>		Most soluble		BaSO <sub>4</sub>		Least soluble

\_\_\_\_\_ charged negative ions tend to \_\_\_\_\_ in solubility as you go down the Group

\_\_\_\_\_ charged negative ions tend to \_\_\_\_\_ in solubility as you go down the Group

**Mg(OH)<sub>2</sub> is said to be sparingly soluble**


**BaSO<sub>4</sub> is insoluble**

## 1) Hydroxides:

- As **solubility increases**, more  $\text{OH}^-$  ions are released.
- This makes a **more alkaline** solution.
- The **pH increases** down the Group

### Testing the solubility of the Group 2 hydroxides:

- This is done by **adding** hydroxide ions,  $\text{OH}^-$  to a solution of the Group 2 ion,  $\text{M}^{2+}$ :




- As  $\text{Mg}(\text{OH})_2$  is sparingly soluble, a thick white precipitate is formed.
- As  $\text{Ba}(\text{OH})_2$  is more soluble, a thin white precipitate will be formed.

## 2) Sulphates:

- Most are actually soluble ranging from sparingly soluble  $\rightarrow$  soluble.
- **Barium sulphate** however is **insoluble**
- This therefore is used as the **chemical test** for the presence of **sulphate ions,  $\text{SO}_4^{2-}$**


### Testing the solubility of the Group 2 sulphates:

- This is done by **adding** sulphate ions,  $\text{SO}_4^{2-}$  or sulphuric acid,  $\text{H}_2\text{SO}_4$  to a solution of the Group 2 ion,  $\text{M}^{2+}$ :




- As  $\text{MgSO}_4$  is soluble, no precipitate will form.
- As  $\text{BaSO}_4$  is insoluble, a white precipitate will be formed.

### Test for sulphate ions, $\text{SO}_4^{2-}$ (Part of Required practical 4)




- Add HCl first – This reacts and removes any sulphites or carbonates that may also give a white precipitate.
- Add  $\text{BaCl}_2$  solution: If sulphates are present, a **white precipitate of  $\text{BaSO}_4$**  will form.



## Test for Group 2 metal ions - Flame tests (Part of Required practical 4)

Nichrome wire dipped in concentrated HCl (to clean) then the unknown compound



<i>Metal ion</i>	<i>Flame colour</i>
<b>Calcium, Ca<sup>2+</sup></b>	<b>Brick red</b>
<b>Strontium, Sr<sup>2+</sup></b>	<b>Red</b>
<b>Barium, Ba<sup>2+</sup></b>	<b>Pale green</b>


### Uses of Group 2 compounds:

#### 1) Barium meals – X – Rays:

- Barium sulphate, BaSO<sub>4</sub>, does not allow X – rays to pass through.
- Drinking a **suspension** of BaSO<sub>4</sub> coats the oesophagus, stomach or intestines (Barium meal)
- These now show up on an X – Ray allowing you to see any problems.
- Other Barium compounds are poisonous.
- Other Group 2 metal compounds are soluble so cannot be used.

#### 2) Extraction of Titanium:


- TiO<sub>2</sub> is converted to TiCl<sub>4</sub> by heating with carbon and chlorine
- TiCl<sub>4</sub> is then reduced by Mg:



- Ti has been **reduced** from **+4 → 0**
- Mg has been **oxidised** from **0 → +2**
- This makes magnesium a **reducing agent**

#### 3) Removal of SO<sub>2</sub> from flue gases:

- SO<sub>2</sub> is produced burning fossil fuels to make electricity.
- This can be removed by reacting with an alkali such as CaO or CaCO<sub>3</sub> slurry (mixed with water)
- The process is called **wet scrubbing**:



#### 4) Neutralising acids:

- Group 2 hydroxides are alkaline and therefore can be used to neutralise acids.
- Ca(OH)<sub>2</sub> is used to neutralise acidic soils.
- Mg(OH)<sub>2</sub> is used to neutralise excess stomach acids.