
1.10 Equilibria and Kp

AS review

- Consider the reaction:



Characteristics of the dynamic equilibrium

- 1) The rate of the forward reaction is equal to the rate of the reverse reaction
- 2) The concentration of the reactants and products are unchanged under stable conditions

Le Chatelier's Principle

When a reaction at equilibrium is subject to a change in concentration, pressure or temperature, the position of the equilibrium will move to counteract the change.

1) Changing concentration – Position of equilibria shifts, K_c unchanged



Increasing concentration:


- **Adding H_2 – increases the concentration of H_2**
- **Equilibrium shifts to the products**
- **To reduce the concentration of the H_2 – counteracting the change**
- The mixture will become less purple

Decreasing concentration:

- **Removing I_2 – decreases the concentration of I_2**
- **Equilibrium shifts to the reactants**
- **To increase the concentration of the I_2 – counteracting the change**
- The mixture will become more purple

The equilibrium moves to oppose the change in concentration

2) Changing pressure – gases only - Position of equilibria shifts, K_c unchanged



Increasing pressure:


- **Equilibrium** shifts to the **reactants**
- This is the side with **fewer moles of gas**
- This will **reduce the pressure – counteracting the change**
- The mixture will become less brown

Decreasing pressure:

- **Equilibrium** shifts to the **products**
- This is the side with **more moles of gas**
- This will **increase the pressure – counteracting the change**
- The mixture will become more brown

The equilibrium moves to oppose the change in pressure

3) Changing temperature - Position of equilibria shifts, K_c changes



Increasing temperature:


- **Equilibrium** shifts to the **products**
- As this is the **endothermic direction**
- This will **decrease temperature – counteracting the change**
- The mixture will become more brown

Decreasing temperature:

- **Equilibrium** shifts to the **reactants**
- As this is the **exothermic direction**
- This will **increase temperature – counteracting the change**
- The mixture will become less brown


The equilibrium moves to oppose the change in temperature

4) The effect of a catalyst on an equilibrium




- A catalyst has **no effect** on the **position of the equilibrium**.
- A catalyst **speeds up the forward and reverse reaction** so it will only increase the rate at which equilibrium is achieved.

The equilibrium constant, K_c




Writing expressions for K_c :

- At equilibrium the concentrations of NO_2 and N_2O_4 are constant:



What is the significance of a K_c


- K_c is a mathematical representation of the ratio of **products : reactants**.






K_p and equilibria:

- K_c and K_p are both expressions to show the position of an equilibria
- c stands for an equilibrium where the individual species are measured using concentrations
- p stands for an equilibrium where the individual species are measured using partial pressures:

Imagine a mixture of gases at equilibrium with a total pressure, $P_t = 100$ KPa



- A total of 20 moles are responsible for the 100 KPa of pressure.
- If we separate the mixture into individual species:

		
10 / 20 th's moles of the filled red particles are responsible for a partial amount of the pressure.	6 / 20 th's moles of the horizontal blue striped particles are responsible for a partial amount of the pressure.	4 / 20 th's moles of the vertical green striped particles are responsible for a partial amount of the pressure.
Partial pressure 'r', P_r	Partial pressure 'b', P_b	Partial pressure 'g', P_g
$P_r = 50$ KPa	$P_b = 30$ KPa	$P_g = 20$ KPa


The partial pressures, p are essentially proportional to the concentrations

The total pressure, P_t is equal to the sum of all of the partial pressures:

$$P_t = P_r + P_b + P_g$$


$$100 = 50 + 30 + 20$$

Writing K_p



- The units of K_p will depend upon the units that pressure is measured in:
Pa KPa mmHg Atm MPa
- It will also depend on the molar ratio in the equilibria equation.

Example:



$$K_p = \frac{P_{\text{products}}}{P_{\text{reactants}}}$$

$$K_p = \frac{P_{\text{SO}_2}^2 \times P_{\text{O}_2}}{P_{\text{SO}_3}^2}$$

Units:

$$K_p = \frac{\text{Pa}^2 \times \text{Pa}}{\text{Pa}^2}$$

$$K_p = \frac{\text{Pa}^2 \times \text{Pa}}{\text{Pa}^2}$$

$$K_p = \text{Pa}$$


Questions:

Write K_p expressions for the following equilibria, for each, work out the units



The equilibrium position and K_p

1) Changing pressure – gases only - Position of equilibria shifts, K_c unchanged



Increasing pressure:

a) Equilibrium position moves to left:

- Equilibrium shifts to the reactants
- This is the side with fewer moles of gas
- This will reduce the pressure – counteracting the change
- The mixture will become less brown

b) K_p is unchanged:



Equilibria shifts to LHS


$$K_p = \frac{P_{\text{NO}_2}^2}{P_{\text{N}_2\text{O}_4}}$$

The increase in pressure also increases the partial pressure of NO_2

Partial pressure of N_2O_4 increases

This maintains K_p

- With an increase in pressure **both partial pressures increase** (smaller volume)
- Equilibrium shifts to the side with fewer moles of gas (LHS) to relieve pressure.
- This will **increase** the **partial pressure** further of the N_2O_4
- The partial pressure of NO_2 decreases slightly due to the shift in equilibrium. The initial increase in the partial pressures at the start outweighs this slight decrease resulting in an overall **increase partial pressure of NO_2**



Decreasing pressure:

a) Equilibrium position moves to right:

- **Equilibrium** shifts to the **products**
- This is the side with **more moles of gas**
- This will **increase the pressure** – **counteracting the change**
- The mixture will become more brown

b) K_p is unchanged:



Equilibria shifts to RHS

$$K_p = \frac{P_{\text{NO}_2}^2}{P_{\text{N}_2\text{O}_4}}$$

The decrease in pressure also decreases the partial pressure of NO_2

Partial pressure of N_2O_4 decreases

This maintains K_p

- With a decrease in pressure **both partial pressures decrease** (larger volume)
- Equilibrium shifts to the side with more moles of gas (RHS) to increase pressure.
- This will **decrease the partial pressure** further of the **N_2O_4**
- The partial pressure of NO_2 increases slightly due to the shift in equilibrium. The initial decrease in the partial pressures at the start outweighs this slight increase resulting in an overall **decrease partial pressure of NO_2**

2) Changing temperature - Position of equilibria shifts, K_c changes



Increasing temperature:

a) Equilibrium shifts to the right hand side

- As this is the **endothermic direction**
- This will **decrease temperature** – counteracting the change
- The mixture will become more brown

b) K_p is increases:




Equilibria shifts to RHS

$$K_p = \frac{P_{\text{NO}_2}^2}{P_{\text{N}_2\text{O}_4}}$$

K_p Increases

This increases the partial pressure of NO_2
 This decreases the partial pressure of N_2O_4

- As equilibria shifts to the products, **P_{NO_2} increases**
- As equilibria shifts to the products, **$P_{\text{N}_2\text{O}_4}$ decreases**
- Products increase / reactants decrease, therefore **K_p increases**



Decreasing temperature:

a) **Equilibrium** shifts to the **reactants**

- As this is the **exothermic direction**
- This will **increase temperature** – counteracting the change
- The mixture will become less brown

b) **K_p** is decreases:



Equilibria shifts to LHS

$$K_p = \frac{P_{\text{NO}_2}^2}{P_{\text{N}_2\text{O}_4}}$$


This decreases the partial pressure of NO₂

This increases the partial pressure of N₂O₄

K_p Decreases

- As equilibria shifts to the products, **P NO₂ decreases**
- As equilibria shifts to the products, **P N₂O₄ increases**
- Products decrease / reactants increase, therefore **K_p decreases**

3) The effect of a catalyst on an equilibrium






- A catalyst has **no effect** on the **position of the equilibrium**.
- A catalyst **speeds up the forward and reverse reaction** so it will only increase the rate at which equilibrium is achieved.

How are partial pressures calculated:

- Remember, each individual species exerts their own partial pressure, which, when all added together gives the total pressure.
- The partial pressure is due to the mole fraction of the total number of moles of that species present in the mixture:

$$\text{Partial pressure} = \text{mole fraction} \times \text{total pressure}$$

$$\text{Mole fraction} = \frac{\text{number of moles of gas}}{\text{total number of moles of gas}}$$

		
Mole fraction: $mf = n / n_t$ $= 10 / 20$ $= 0.5$ Partial pressure: $P_r = mf \times P_t$ $= 0.5 \times 100$ $= 50$	Mole fraction: $mf = n / n_t$ $= 6 / 20$ $= 0.3$ Partial pressure: $P_r = mf \times P_t$ $= 0.3 \times 100$ $= 30$	Mole fraction: $mf = n / n_t$ $= 4 / 20$ $= 0.2$ Partial pressure: $P_r = mf \times P_t$ $= 0.2 \times 100$ $= 20$
$P_r = 50 \text{ KPa}$	$P_b = 30 \text{ KPa}$	$P_g = 20 \text{ KPa}$



$$K_p = \frac{P_b \times P_g}{P_r}$$

$$K_p = \frac{30 \times 20}{50}$$

$$K_p = 1.2 \text{ KPa}$$

Example:

The following equilibria was found to contain 3 moles of HI, 6 moles of H₂ and 2 moles of I₂. The total pressure of the sealed vessel was 100KPa

	2HI	\rightleftharpoons	H ₂	+	I ₂
Moles of gas:	3		6		1
Mole fraction:	3/10		6/10		1/10
Partial pressure:	3/10 x 100		6/10 x 100		1/10 x 100
Partial pressure:	30		60		10

$$K_p = \frac{P_{H_2} \times P_{I_2}}{P_{HI}^2}$$


$$K_p = \frac{60 \times 10}{(30)^2}$$

$$K_p = \frac{600}{900}$$

$$K_p = 0.67 \quad \text{No units (cancel out)}$$

Questions: Give all answers to 3SF

- 1) The following equilibria was found to contain 5 moles of HCl, 10 moles of H₂ and 8 moles of Cl₂. The total pressure of the sealed vessel was 50KPa



- Write an expression for K_p
- Calculate the mole fractions for each gas
- Calculate the partial pressures for each gas
- Calculate K_p for the equilibria. Include units in your answer.

Ans = 3.18

- 2) The following equilibria was found to contain 5 moles of NO₂ and 8 moles of N₂O₄. The total pressure of the sealed vessel was 760 mmHg



Calculate K_p. Include units and show all working out clearly:

Ans = 183

- 3) The following equilibria was found to contain 5 moles of SO₂, 5 moles of O₂ and 8 moles of SO₃. The total pressure of the sealed vessel was 2 atm



Calculate K_p. Include units and show all working out clearly:

Ans = 4.59

- 4) An 80 dm³ reaction vessel contains a mixture of hydrogen and oxygen with a total pressure is 300 kPa. The partial pressure of oxygen 20 kPa.

a) Calculate the partial pressure of hydrogen.

b) Calculate the mole fraction of hydrogen and oxygen.

- 5) The following equilibria was established. It has a K_p of 50 and the partial pressures of H_2 and I_2 are 250 and 160 respectively.




- a) Write an expression for K_p

- b) Rearrange the expression and calculate the partial pressure of HI

Ans = 28.3

- 6) The following equilibria was established. It has a K_p of $7.48 \times 10^{-8} \text{ kPa}^{-2}$. The mixture has a total pressure of 5000KPa. The partial pressures of H_2 and NH_3 are 2814 and 1250 respectively.



Calculate the mole fraction of N_2

Ans = 0.187

Challenging:

1) 3.00 moles of $\text{SO}_3(\text{g})$ are placed into an 8.00 dm^3 container and heated to 1105 K.



- At equilibrium the mixture contains 0.58 mol of $\text{O}_2(\text{g})$.
- The total pressure of the equilibrium mixture is $3.45 \times 10^6 \text{ Pa}$.
- Use this information to calculate K_p

a) Determine equilibrium moles of all gases.

b) Determine the mole fractions of each of the gases.

c) Calculate the partial pressures.

d) Write the expression for K_p .

e) Calculate K_p and state its units.

Ans = 222000

2) 9.20 g of $\text{N}_2\text{O}_4(\text{g})$ was heated to a temperature of 340 K at a pressure of 13.3 kPa.



Once equilibrium had been reached, 70% of the $\text{N}_2\text{O}_4(\text{g})$ had dissociated.

Use this information to calculate K_p

Ans = 51.5